

Human Recombinant IFN-gamma, ACF

Interferon-gamma, animal component-free

Catalog #78141	20 µg
Catalog #78141.1	100 µg
Catalog #78141.2	500 µg
Catalog #78141.3	1000 µg

Product Description

Interferon-gamma (IFN- γ), also known as type II interferon, is produced by T and NK cells, and in smaller amounts by dendritic cells and macrophages. IFN- γ is controlled by cytokines such as IL-12 and IL-18 secreted in response to infection (Schroder et al.). IFN- γ binds to a receptor complex and initiates signal transduction via the JAK/STAT pathway; this culminates in the transcription and activation of many genes that control a diverse array of immunological functions (de Weerd and Nguyen; Krause et al.). IFN- γ stimulates the antimicrobial and anti-tumor activity of macrophages, NK cells, and neutrophils (Billiau & Matthys) by promoting the activation of microbial effector functions such as production of reactive oxygen species, nitric oxide intermediates, and complement (Schroder et al.). IFN- γ enhances the expression of major histocompatibility complex class I and II in dendritic cells and mononuclear phagocytes, as well as the production of IL-12 by dendritic cells. In B cells, IFN- γ stimulates survival and growth in both mouse and human cells, and redirects B cells from proliferation towards differentiation. IFN- γ favors the development of Th1 vs Th2 cells and stimulates monocyte differentiation and function (Schroder et al.). This product is animal component-free.

Product Information

Alternative Names:	Interferon gamma, Type II interferon
Accession Number:	P01579
Amino Acid Sequence:	MQDPYVKEAE NLKKYFNAGH SDVADNGTLF LGILKNWKEE SDRKIMQSQI VSFYFKLFKN FKDDQSIQKS VETIKEDMNV KFFNSNKKKR DDFEKLTNYS VTDLNVQRKA IHELIQVMAE LSPA AKTGKR KRSQMLFQGR RASQ
Predicted Molecular Mass:	16.9 kDa
Species:	Human
Product Formulation:	Lyophilized from a sterile-filtered aqueous solution containing sodium phosphate, trehalose, mannitol, and TWEEN® 80, pH 6.5
Source:	E. coli
Purity:	≥ 95%

Specifications

Activity:	The specific activity is $\geq 3.3 \times 10^6$ units/mg (≤ 0.3 ng/mL), as determined by the ability to induce cytotoxicity of HT-29 cells.
Endotoxin Level:	Measured by kinetic Limulus amoebocyte lysate (LAL) analysis and is ≤ 0.1 EU/ μ g protein.

Preparation and Storage

Stability and Storage:	Store at -20 to -80°C. Stable as supplied for 12 months from date of receipt.
Preparation:	<p>Centrifuge vial before opening. Bring vial and sterile water to room temperature (15 - 25°C). Reconstitute the product in sterile water to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. Let solution sit for 1 minute at room temperature. If precipitate is observed, centrifuge at 16,000 x g for 1 minute. Remove supernatant and transfer to a new tube, taking care not to disturb the pellet. Discard the pellet. A 10% overfill has been added to compensate for any loss of protein in the precipitate.</p> <p>OPTIONAL: After reconstitution, if product will not be used immediately, dilute with concentrated bovine serum albumin (BSA) to a final BSA concentration of 0.1%. The effect of storage of stock solution on product performance should be tested for each application. As a general guide, do not store at 2 - 8°C for more than 1 month or at -80°C for more than 3 months. Avoid repeated freeze-thaw cycles.</p>

Data




Figure 1. Biological Activity of Human Recombinant IFN-gamma, ACF

(A) The biological activity of Human Recombinant IFN-gamma, ACF was measured by the ability to induce cytotoxicity of HT-29 cells. The EC₅₀ is defined as the effective concentration of the cytokine at which cytotoxicity is at 50% of maximum. The EC₅₀ in the example above is ≤ 0.3 ng/mL.

(B) 1 μ g of Human Recombinant IFN-gamma, ACF was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human Recombinant IFN-gamma, ACF has a predicted molecular mass of 16.9 kDa.

Related Products

For a complete list of cytokines or peptide pools, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

- Billiau A & Matthys P. (2009) Interferon-gamma: a historical perspective. *Cytokine Growth Factor Rev* 20(2): 97–113.
- de Weerd NA & Nguyen T. (2012) The interferons and their receptors--distribution and regulation. *Immunol Cell Biol* 90(5): 483–91.
- Krause CD et al. (2000) Signaling by covalent heterodimers of interferon-gamma. Evidence for one-sided signaling in the active tetrameric receptor complex. *J Biol Chem* 275(30): 22995–3004.
- Schroder K et al. (2004) Interferon-gamma: an overview of signals, mechanisms and functions. *J Leukoc Biol* 75(2): 163–89.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2025 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. TWEEN is a registered trademark of Croda Americas LLC. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.