

Anti-Human CD90 Antibody, Clone 5E10

Antibodies

Mouse monoclonal IgG1 antibody
against human, rhesus, cynomolgus
CD90 (Thy-1), unconjugated



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog #60045

100 µg 0.5 mg/mL

FOR RESEARCH USE ONLY. NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES.

Product Description

The 5E10 antibody reacts with CD90 (Thy-1), a GPI-linked membrane glycoprotein that is N-glycosylated at two sites, giving rise to 25 - 37 kDa molecules. CD90 has roles in signal transduction, cell adhesion and migration, neurite outgrowth, T cell activation, tumor suppression, and inhibition of the proliferation and differentiation of hematopoietic stem cells. It is a known ligand of $\beta 2$ and $\beta 3$ integrins and upregulates synthesis of fibronectin, osteonectin and thrombospondin. CD90 is broadly expressed, being found on human thymocytes, neurons, some glial cells, fibroblasts, activated endothelial cells, some leukemia cell lines and a distinct subset (<1%) of CD3+CD4+ T cells in human peripheral blood. CD90 is also expressed by small subsets of CD34+ cells in fetal liver, umbilical cord blood, bone marrow and mobilized peripheral blood cells. CD90 is considered an important marker for hematopoietic stem and progenitor cells and, in combination with other markers such as CD34, is useful to identify and isolate these cells by FACS.

Target Antigen Name:	CD90 (Thy-1)
Alternative Names:	Thy-1, Thy1, CDw90
Gene ID:	7070
Species Reactivity:	Human, Rhesus, Cynomolgus, Baboon, Pigtailed Macaque, Dog, Pig
Host Species:	Mouse (BALB/c)
Clonality:	Monoclonal
Clone:	5E10
Isotype:	IgG1, kappa
Immunogen:	Human HEL erythroleukemia cell line
Conjugate:	Unconjugated

Applications


Verified:	FC
Reported:	FC, ICC, IF, IHC, IP, WB
Special Applications:	This antibody clone has been verified for labeling human mesenchymal cells grown in MesenCult™ Proliferation Kit (Human; Catalog #05411) and MesenCult™-XF Medium (Catalog #05420).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; WB: Western blotting

Properties

Formulation:	Phosphate-buffered saline
Purification:	The antibody was purified by affinity chromatography.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. Addition of 0.1% sodium azide (final) is recommended once the vial has been opened. For product expiry date, please contact techsupport@stemcell.com .
Directions for Use:	For flow cytometry the suggested use of this antibody is ≤ 1 µg per 1×10^6 cells in 100 µL volume or per 100 µL of whole blood. It is recommended that the antibody be titrated for optimal performance for each application.

Data



Flow cytometry analysis of human HEL cells labeled with Anti-Human CD90 Antibody, Clone 5E10, followed by goat anti-mouse IgG, FITC (filled histogram). Labeling with a mouse IgG1, kappa isotype control antibody followed by goat anti-mouse IgG, FITC is shown (open histogram).

Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

References

1. Craig W, et al. Expression of Thy-1 on human hematopoietic progenitor cells. J Exp Med 177(5): 1331-42, 1993 (FC, IP, WB)
2. Holden JT, et al. Characterization of Thy-1 (CDw90) expression in CD34+ acute leukemia. Blood 86(1): 60-65, 1995
3. Mayani H, Lansdorp PM. Thy-1 expression is linked to functional properties of primitive hematopoietic progenitor cells from human umbilical cord blood. Blood 83(9): 2410-07, 1994 (FC)
4. Murray LJ, Tsukamoto A, Hoffman R. CD34+Thy-1+Lin- stem cells from mobilized peripheral blood. Leuk Lymphoma 22(1-2): 37-42, 1996 (FC)
5. Mason D, et al. Eds. Leukocyte Typing VII: White Cell Differentiation Antigens. Oxford University Press, Oxford, UK, p. 836, 2002
6. Hung JT, et al. Immunopathogenic role of TH1 cells in autoimmune diabetes: evidence from a T1 and T2 doubly transgenic non-obese diabetic mouse model. J Autoimmun 25(3):181-92, 2005 (IHC, FC)

Copyright © 2013 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists and MesenCult are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485 MEDICAL DEVICE STANDARDS.