

EasySep™ EasySep™ "Do-It-Yourself" Selection Kit

Positive Selection
Catalog #18098

For processing 1 x 10⁹ cells



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713
INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM
FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Document #28863 | Version 2_1_0

Description

Isolate highly purified cell types of interest with your own mouse IgG1 monoclonal antibody from single-cell suspensions by immunomagnetic positive selection.

- Fast and easy-to-use
- No columns

This kit targets any cell type of interest with your own mouse IgG1 monoclonal antibody. Desired cells are labeled with antibodies and magnetic particles, and separated without columns using an EasySep™ magnet. Unwanted cells are simply poured off, while desired cells remain in the tube. Isolated cells are immediately available for downstream applications such as flow cytometry, culture, or DNA/RNA extraction.

Component Descriptions

COMPONENT NAME	COMPONENT #	QUANTITY	STORAGE	SHELF LIFE	FORMAT
EasySep™ "Do-It-Yourself" Component A	18090	1 x 0.1 mL	Store at 2 - 8°C. Do not freeze.	Stable until expiry date (EXP) on label.	A combination of monoclonal antibodies in PBS.
EasySep™ "Do-It-Yourself" Component B	18091	1 x 0.1 mL	Store at 2 - 8°C. Do not freeze.	Stable until expiry date (EXP) on label.	A combination of monoclonal antibodies in PBS.
EasySep™ Magnetic Nanoparticles Positive Selection	18150	1 x 1 mL	Store at 2 - 8°C. Do not freeze.	Stable until expiry date (EXP) on label.	A suspension of magnetic particles in water.
RoboSep™ Vial for "Do-It-Yourself" Antibody Cocktail	18093	1 vial	Not applicable	Not applicable	Not applicable

PBS - phosphate-buffered saline

Components may be shipped at room temperature (15 - 25°C) but should be stored as indicated above.

Additional Reagent Stability Information

REAGENT NAME	STORAGE	SHELF LIFE
Selection Cocktail (your own mouse IgG1 monoclonal antibody + Component A + Component B)	Store at 2 - 8°C. Do not freeze.	Stability should be experimentally determined. Do not exceed the shelf life of the individual components.

Sample Preparation

For available fresh and frozen samples, see www.stemcell.com/primarycells.

PERIPHERAL BLOOD

Prepare a peripheral blood mononuclear cell (PBMC) suspension from whole blood by centrifugation over a density gradient medium (e.g. Lymphoprep™, Catalog #07801). For more rapid PBMC preparation, use the SepMate™ RUO (Catalog #86450/86415) or SepMate™ IVD* (Catalog #85450/85415) cell isolation tube. If using previously frozen PBMCs, incubate the cells with DNase I Solution (Catalog #07900) at a concentration of 100 µg/mL at room temperature (15 - 25°C) for at least 15 minutes prior to labeling and separation. Filter clumpy suspensions through a 40 µm Cell Strainer (Catalog #27305) for optimal results.

After preparation, resuspend cells at 1 x 10⁸ cells/mL in recommended medium.

* SepMate™ IVD is only available in select regions where it is registered as an In Vitro Diagnostic (IVD) device for the isolation of mononuclear cells (MNCs) from whole blood or bone marrow by density gradient centrifugation. In all other regions SepMate™ is available for research use only (RUO).

OTHER SAMPLE SOURCES

If other sources or tissues are used, ensure that the cells are in a single-cell suspension. Not recommended for selection of mouse cells.

Recommended Medium

EasySep™ Buffer (Catalog #20144), RoboSep™ Buffer (Catalog #20104), or PBS containing 2% fetal bovine serum (FBS) and 1 mM EDTA. Medium should be free of Ca⁺⁺ and Mg⁺⁺.

Directions for Use – Manual EasySep™ Protocols

See page 1 for Sample Preparation and Recommended Medium. Refer to Table 1 for detailed instructions regarding the EasySep™ procedure.

Table 1. EasySep™ "Do-It-Yourself" Selection Kit Protocol

STEP	INSTRUCTIONS	 EasySep™ (Catalog #18000)
1	Add your own mouse IgG1 monoclonal antibody (reconstituted if necessary) to a 1.5 mL polypropylene tube.	15 µg NOTE: Up to 800 µL
2	Add Component A to the polypropylene tube and mix well.	100 µL
3	Add Component B to the polypropylene tube.	100 µL
	Mix and incubate.	37°C for 5 hours or overnight
4	Add sterile PBS to top up the polypropylene tube to the indicated volume.	Top up to 1 mL
Prepared Selection Cocktail is ready to use.		---
5	Prepare sample at the indicated cell concentration within the volume range.	1 x 10 ⁸ cells/mL 0.1 - 2.5 mL NOTE: If starting with fewer than 1 x 10 ⁷ cells, resuspend cells in 0.1 mL For samples with a starting frequency of desired cells < 10%, start with a concentration of 2 x 10 ⁸ cells/mL
	Add sample to required tube.	5 mL (12 x 75 mm) polystyrene round-bottom tube (e.g. Corning Catalog #352058)
6	Add species-specific FcR blocker (not provided) to sample (if desired).	0.5 - 3 µg/mL of sample
7	Add Selection Cocktail to sample.	100 µL/mL of sample
	Mix and incubate.	RT for 15 minutes
8	Mix Magnetic Particles. NOTE: Particles should appear evenly dispersed.	Pipette up and down more than 5 times
9	Add Magnetic Particles to sample.	50 µL/mL of sample
	Mix and incubate.	RT for 10 minutes
10	Add recommended medium to top up the sample to the indicated volume. Mix by gently pipetting up and down 2 - 3 times.	Top up to 2.5 mL
	Place the tube (without lid) into the magnet and incubate.	RT for 5 minutes*
11	Pick up the magnet, and in one continuous motion invert the magnet and tube,** pouring off the supernatant. Remove the tube from the magnet; this tube contains the isolated cells.	Discard supernatant
12	Repeat steps as indicated.	Steps 10 and 11, two more times (total of 3 x 5-minute separations)
OPTIONAL ADDITIONAL SEPARATION(S) For samples with a starting frequency of desired cells < 10% NOTE: This will improve purity but may reduce recovery		Repeat steps 10 and 11, up to three more times (total of 4 - 6 x 5-minute separations)
13	Resuspend cells in desired medium. Be sure to collect cells from the sides of the tube.	Isolated cells are ready for use

RT - room temperature (15 - 25°C)

* Recovery may be improved by increasing separation time in the magnet to 10 minutes for each round.

** Leave the magnet and tube inverted for 2 - 3 seconds, then return upright. Do not shake or blot off any drops that may remain hanging from the mouth of the tube.

Directions for Use – Fully Automated RoboSep™ Protocol

See page 1 for Sample Preparation and Recommended Medium. Refer to Table 2 for detailed instructions regarding the RoboSep™ procedure.

Table 2. RoboSep™ "Do-It-Yourself" Selection Kit Protocol

STEP	INSTRUCTIONS	RoboSep™ (Catalog #20000 and #21000)	
1	Add your own mouse IgG1 monoclonal antibody (reconstituted if necessary) to the RoboSep™ Vial for "Do-It-Yourself" Antibody Cocktail provided.	15 µg NOTE: Up to 800 µL	
2	Add Component A to the RoboSep™ Vial and mix well.	100 µL	
3	Add Component B to the RoboSep™ Vial.	100 µL	
	Mix and incubate.	37°C for 5 hours or overnight	
4	Add sterile PBS to top up the RoboSep™ Vial to the indicated volume.	Top up to 1 mL	
Prepared Selection Cocktail is ready to use.		---	
5	Prepare sample at the indicated cell concentration within the volume range.	1 x 10 ⁸ cells/mL 0.25 - 8.5 mL NOTE: For samples with a starting frequency of desired cells < 10%, start with a concentration of 2 x 10 ⁸ cells/mL	
	Add sample to required tube.	14 mL (17 x 100 mm) polystyrene round-bottom tube (e.g. Corning Catalog #352057)	
6	Add species-specific FcR blocker (not provided) to sample (if desired).	0.5 - 3 µg/mL of sample	
7	Select protocol.	Any Species Do-It-Yourself Positive Selection 18098-high purity	
8	Mix Magnetic Particles. NOTE: Particles should appear evenly dispersed.	Pipette up and down more than 5 times	
9	Load the carousel.	Follow on-screen prompts	
	Start the protocol.	Press the green "Run" button	
10	Unload the carousel when the run is complete. Remove the tube containing the isolated cells and resuspend in desired medium. Be sure to collect cells from the sides of the tube.	Isolated cells are ready for use	

Notes and Tips

ASSESSING PURITY

The mouse IgG1 antibody used in the cocktail created with the EasySep™ "Do-It-Yourself" Selection Kit may block other fluorochrome-conjugated antibodies that are used to assess purity by flow cytometry. Use fluorochrome-conjugated antibody clones that are not blocked by the antibody clone used in the Selection Cocktail. If this is not possible, one of the following methods can be used to assess purity:

- Use alternative fluorochrome-conjugated markers, if applicable.
- Use a fluorochrome-conjugated secondary antibody, such as Goat Anti-Mouse IgG (H+L) Antibody, Polyclonal (Catalog #60138).

ANTIBODY QUALITY

Recovery of positively selected cells is dependent on the quality of the mouse IgG1 antibody used in the cocktail created with the EasySep™ "Do-It-Yourself" Selection Kit. Antibodies that have expired or that have been stored improperly may show lower affinity for the surface marker on the target cell, resulting in lower recovery.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, EasySep, RoboSep, and SepMate are trademarks of STEMCELL Technologies Inc. Lymphoprep is a trademark of AXIS-SHIELD. All other trademarks are the property of their respective holders. Users of this selection kit should ensure that they are entitled to use the antibody of interest. STEMCELL Technologies is not responsible for patent infringements or violations that may occur when using this product. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.