

PRODUCT DESCRIPTION:

A combination of mouse and rat monoclonal antibodies purified from mouse ascites or hybridoma culture supernatant. Purified by affinity chromatography using Protein A or Protein G Sepharose. These antibodies are bound in bispecific Tetrameric Antibody Complexes (TAC), which are directed against HLA-DR and dextran. The subclass of the mouse monoclonal antibody is IgG₁.

SPECIFICITY:

This TAC recognizes human HLA-DR cell surface antigens and dextran. HLA-DR is a major histocompatibility class II antigen expressed on B cells, activated T cells, and antigen presenting cells.

FORMAT:

Catalog #14236 – contains 1 x 14236C

Catalog #14256 – contains 2 x 14236C

Supplied at a concentration of **300 µg/mL** in phosphate buffered saline with trace amounts of borate (6 mM). Does not contain sodium azide or BSA.

It should be kept in mind that this product is a biological reagent, and as such cannot be completely characterized or quantified. Some variability is unavoidable.

STABILITY AND STORAGE:

Store at 4°C. DO NOT FREEZE. Stable for 2 years.
Contents sterile in unopened tube.

APPLICATIONS:

Deplete HLA-DR⁺ (MHC class II⁺) cells from human peripheral blood mononuclear cell samples by combining HLA-DR TAC with StemSep[®] or EasySep[®] magnetic cell separation.

To isolate resting human CD4⁺ or CD8⁺ T cells:

Add HLA-DR TAC with either **StemSep[®] CD4⁺** or **CD8⁺ T Cell Enrichment** cocktails (Catalog #14052, 14053) or **EasySep[®] CD4⁺** or **CD8⁺ T Cell Enrichment** cocktails (Catalog #19052, 19053).

DIRECTIONS FOR USE:

Centrifuge tube before using to ensure recovery of entire contents.

Add TAC at 10 µL/mL of cells prepared at a concentration of 5×10^7 cells/mL. Titration in the range of **0.1 – 3.0 µg/mL** final concentration may be required for optimal performance. When adding TAC with an enrichment cocktail, add the TAC during the same step as adding the cocktail to the cells.

Please contact us for detailed protocol information.
Refer to Material Safety Data Sheet for more information.

REFERENCES:

1. Lansdorp PM, Aalberse RC, Bos R, Schutter W and Van Bruggen EFJ. Cyclic tetramolecular complexes of monoclonal antibodies: A new type of cross-linking reagent. Eur. J. Immunol. 1986; 16: 679.
2. Thomas TE, Sutherland HJ and Lansdorp PM. Specific binding and release of cells from beads using cleavable tetrameric antibody complexes. J. Immunol. Methods 1989; 120: 221.
3. Chun T-W, Engel D, Berrey MM, Shea T, Corey L, Fauci AS: Early establishment of a pool of latently infected, resting CD4⁺ T cells during primary HIV-1 infection. Proc Natl Acad Sci USA 1998; 95: 8869.

**THIS REAGENT IS FOR RESEARCH ONLY.
IT IS NOT TO BE ADMINISTERED TO HUMANS.**

StemCell Technologies

In North America

Tel: 1.604.877.0713
Fax: 1.604.877.0704
Toll Free Tel: 1.800.667.0322
Toll Free Fax: 1.800.567.2899
e-mail: info@stemcell.com
www.stemcell.com

In the United Kingdom

Tel: +44.(0)20.7537.7565
Fax: +44.(0)20.7515.5408
Toll Free within United Kingdom:
Tel: 0800.731.27.14
Fax: 0800.731.27.13
e-mail: info@stemcellgb.com

In Europe

Tel: +33.(0)4.76.04.75.30
Fax: +33.(0)4.76.18.99.63
e-mail: info@stemcellfrance.com

Revised:
November 2006

Product Information Sheet



Version 2.0.0

ANTI-HUMAN HLA-DR TAC

Tetrameric Antibody
Complex

Catalog #14236

Catalog #14256

100 µL

200 µL