# Anti-Mouse CD11b Antibody, Clone M1/70, APC

### **Antibodies**

Rat monoclonal IgG2b antibody against human, mouse, rhesus CD11b,

APC-conjugated

Catalog #60001AZ #60001AZ.1 100 μg 0.2 mg/mL 25 μg 0.2 mg/mL



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

## **Product Description**

The M1/70 antibody reacts with CD11b, an ~170 kDa type 1 transmembrane glycoprotein which associates non-covalently with CD18 to form the heterodimeric Mac-1 receptor. Through its interactions with ligands such as ICAM-1 (CD54), ICAM-2 (CD102), ICAM-4 (CD242), iC3b, and fibrinogen, Mac-1 functions in several processes, including the adherence of neutrophils and monocytes to stimulated endothelium and phagocytosis of complement-coated particles. CD11b is expressed on the surface of granulocytes, monocytes, NK cells, dendritic cells, tissue macrophages and subsets of T and B cells, and has been used as a marker to distinguish naïve and memory CD8+ T cells. CD11b is a relatively late marker for myeloid differentiation and is undetectable on most myelomonocytic hematopoietic progenitor cells and more primitive cells. The M1/70 antibody reportedly blocks iC3b binding to Mac-1.

Target Antigen Name: CD11b

Alternative Names: alphaM integrin, C3biR, CR3, Ly-40, Mac-1, Mo1

Gene ID: 16409

Species Reactivity: Human, Mouse, Rhesus, Cynomolgus, Baboon, Chimpanzee, Rabbit

Host Species: Rat

Clonality: Monoclonal Clone: M1/70

Isotype: IgG2b, kappa

Immunogen: C57BL/10 mouse splenocytes

Conjugate: APC

# **Applications**

Verified: FC Reported: FC

Special Applications: This antibody clone has been verified for purity assessments of cells isolated with EasySep™ kits, including

EasySep™ Mouse Monocyte Enrichment Kit (Catalog #19761).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

# **Properties**

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide

Purification: The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The

solution is free of unconjugated APC and unconjugated antibody.

Stability and Storage: Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to

light. For product expiry date, please contact techsupport@stemcell.com.

Directions for Use: For flow cytometry the suggested use of this antibody is  $\leq 0.25 \,\mu g$  per 1 x 10<sup>6</sup> cells in 100  $\mu L$  volume. It is


recommended that the antibody be titrated for optimal performance for each application.

#### Anti-Mouse CD11b Antibody, Clone M1/70, APC

## **Antibodies**



### Data



- (A) Flow cytometry analysis of C57BL/6 mouse splenocytes labeled with Anti-Mouse CD11b Antibody, Clone M1/70, APC and Anti-Mouse CD45 Antibody, Clone 30-F11, PE (Catalog #60030PE).
- (B) Flow cytometry analysis of C57BL/6 mouse splenocytes labeled with Rat IgG2b, kappa Isotype Control Antibody, Clone RTK4530, APC (Catalog #60077AZ) and Anti-Mouse CD45 Antibody, Clone 30-F11, PE.

### Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

### References

- 1. Ahn G-O et al. (2010) Inhibition of Mac-1 (CD11b/CD18) enhances tumor response to radiation by reducing myeloid cell recruitment. Proc Natl Acad Sci USA 107(18): 8363–8. (FA, ICC, IF, IHC)
- 2. Baumgartner CK et al. (2010) Peptide-MHC class II complex stability governs CD4 T cell clonal selection. J Immunol 184(2): 573-81. (FACS, FC)
- 3. Charles N et al. (2010) Basophils and the T helper 2 environment can promote the development of lupus nephritis. Nat Med 16(6): 701-7. (FC)
- 4. Norian LA et al. (2009) Tumor-infiltrating regulatory dendritic cells inhibit CD8+ T cell function via L-arginine metabolism. Cancer Res 69(7): 3086–94. (FC)
- 5. Dzhagalov I et al. (2007) The antiapoptotic protein McI-1 is essential for the survival of neutrophils but not macrophages. Blood 109(4): 1620-6. (FC)
- 6. Tailleux L et al. (2003) DC-SIGN is the major Mycobacterium tuberculosis receptor on human dendritic cells. J Exp Med 197(1): 121–7. (FA, FC)
- 7. Iwasaki A & Kelsall BL. (2001) Unique functions of CD11b+, CD8 alpha+, and double-negative Peyer's patch dendritic cells. J Immunol 166(8): 4884–90. (FACS, FC, IF, IHC)
- 8. Noel GJ et al. (1990) Role of complement in mouse macrophage binding of Haemophilus influenzae type b. J Clin Invest 85(1): 208-18. (FA)
- 9. Sanchez-Madrid F et al. (1983) Mapping of antigenic and functional epitopes on the alpha- and beta-subunits of two related mouse glycoproteins involved in cell interactions, LFA-1 and Mac-1. J Exp Med 158(2): 586–602. (IP)
- 10. Beller DI et al. (1982) Anti-Mac-1 selectively inhibits the mouse and human type three complement receptor. J Exp Med 156(4): 1000-9. (FA)
- 11. Ault KA & Springer TA. (1981) Cross-reaction of a rat-anti-mouse phagocyte-specific monoclonal antibody (anti-Mac-1) with human monocytes and natural killer cells. J Immunol 126(1): 359–64. (FA, FACS, FC, RIA)
- 12. Springer T et al. (1979) Mac-1: a macrophage differentiation antigen identified by monoclonal antibody. Eur J Immunol 9(4): 301-6.
- 13. Springer T et al. (1978) Monoclonal xenogeneic antibodies to murine cell surface antigens: identification of novel leukocyte differentiation antigens. Eur J Immunol 8(8): 539–51. (IP)

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists and EasySep are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. Alexa Fluor® is a registered trademark of Life Technologies Corporation. This product is licensed for internal research use only and its sale is expressly conditioned on the buyer not using it for manufacturing, performing a service, or medical test, or otherwise generating revenue. For use other than research, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. While STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.