

Small Molecules

GSA 10

Hedgehog pathway activator; SMO activator

Catalog # 73172
73174

10 mg
50 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM


FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

FOR RESEARCH USE ONLY. NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES.

Product Description

GSA 10 is an agonist of smoothened (SMO), a cell surface receptor and mediator of the hedgehog signaling pathway. It is a quinolinecarboxamide derivative that binds in a distinct binding pocket from cyclopamine. GSA10 is active at an EC_{50} of 1.2 μ M in an alkaline phosphatase-based mouse mesenchymal cell line (C3H10T1/2) differentiation assay (Gorojankina et al.).

| | |
|--------------------|---|
| Molecular Name: | GSA 10 |
| Alternative Names: | Not applicable |
| CAS Number: | 300833-95-8 |
| Chemical Formula: | $C_{26}H_{30}N_2O_5$ |
| Molecular Weight: | 450.5 g/mol |
| Purity: | $\geq 95\%$ |
| Chemical Name: | 4-[[[(1-hexyl-1,2-dihydro-4-hydroxy-2-oxo-3-quinolinyl)carbonyl]amino]-benzoic acid, propyl ester |
| Structure: | |



Properties

| | |
|----------------------|--|
| Physical Appearance: | A crystalline solid |
| Storage: | Product stable at -20°C as supplied. Protect from prolonged exposure to light. For product expiry date, please contact techsupport@stemcell.com. |
| Solubility: | · DMF ≤ 2 mM For example, to prepare a 1 mM stock solution in DMF, resuspend 10 mg in 22.2 mL of DMF. |

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMF at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMF concentration above 0.1% due to potential cell toxicity.

Published Applications

DIFFERENTIATION

- Promotes differentiation of multipotent mesenchymal progenitor cells into osteoblasts (Gorojankina et al.).

References

Gorojankina T et al. (2013) Discovery, molecular and pharmacological characterization of GSA-10, a novel small-molecule positive modulator of Smoothened. Mol Pharmacol 83(5): 1020–9.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, please visit our website at www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

Copyright © 2015 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.