QNZ

Small Molecules

NF-kB inhibitor

Catalog # 73352 1 mg



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

FOR RESEARCH USE ONLY. NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES.

## **Product Description**

QNZ is a quinazoline derivative that inhibits nuclear factor (NF)- $\kappa$ B activation (IC<sub>50</sub> = 11 nM in human Jurkat T lymphocyte cells). NF- $\kappa$ B enhances the transcription of pro-inflammatory cytokines, and QNZ inhibits lipopolysaccharide (LPS)-stimulated tumor necrosis factor (TNF)- $\alpha$  production in mouse splenocytes (IC<sub>50</sub> = 7 nM;Tobe et al.), as well as CXCL1-mediated pro-inflammatory increase in potassium currents in adult rat neurons (Yang et al). It does not inhibit kinases in a standard screen (Wu et al.).

Molecular Name: QNZ

Alternative Names: CAY10470 
CAS Number: 545380-34-5 
Chemical Formula:  $C_{22}H_{20}N_4O$  
Molecular Weight: 356.4 g/mol 
Purity:  $\geq$  98%

Chemical Name: N4-[2-(4-phenoxyphenyl)ethyl]-4,6-quinazolinediamine

Structure:

# **Properties**

Physical Appearance: A crystalline solid

Storage: Product stable at -20°C as supplied. Protect from prolonged exposure to light. For product expiry date, please

contact techsupport@stemcell.com.

**Solubility:**  $\cdot$  DMSO  $\leq$  55 mM

· Absolute ethanol ≤ 25 mM

For example, to prepare a 10 mM stock solution in DMSO, resuspend 1 mg in 281 µL of DMSO.

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

# Small Molecules QNZ



## **Published Applications**

#### DISEASE MODELING

· Blocks amyloid precursor protein release in human SH-SY5Y neuroblastoma cells caused by muscarinic receptor activation (Choi et al.).

#### **MAINTENANCE**

· Neuroprotective in a glutamate toxicity assay using YAC128 medium spiny neuron cultures (Wu et al.).

### References

Choi S et al. (2006) Nuclear factor-kappaB activated by capacitative Ca2+ entry enhances muscarinic receptor-mediated soluble amyloid precursor protein (sAPPalpha) release in SH-SY5Y cells. J Biol Chem 281(18): 12722–8.

Tobe M et al. (2003) Discovery of quinazolines as a novel structural class of potent inhibitors of NF-kappa B activation. Bioorg Med Chem 11(3): 383–91.

Wu J et al. (2011) Neuronal store-operated calcium entry pathway as a novel therapeutic target for Huntington's disease treatment. Chem Biol 18(6): 777–93.

Yang R-H et al. (2009) NF-kappaB mediated enhancement of potassium currents by the chemokine CXCL1/growth related oncogene in small diameter rat sensory neurons. Mol Pain 5:26.

### Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, please visit our website at www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

Copyright © 2015 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.