Anti-Mouse CD19 Antibody, Clone 1D3, FITC

Antibodies

Rat monoclonal IgG2a antibody against mouse CD19, FITC-conjugated

Catalog #60112FI 500 μg 0.5 mg/mL #60112FI.1 100 μg 0.5 mg/mL #60112FI.2 25 μg 0.5 mg/mL



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1800 667 0322 • PHONE +1604 877 0713
INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM
FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

FOR RESEARCH USE ONLY NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPFUTIC USES.

Product Description

The 1D3 antibody reacts with mouse CD19, an ~95 kDa type 1 transmembrane glycoprotein expressed on the surface of B cells throughout all stages of development, from early pre-B cells to plasma cells. Expression is down-regulated but persists in terminally differentiated plasma cells. CD19 is also found on follicular dendritic cells. By associating with CD21 and CD81, CD19 functions as a coreceptor for the B cell receptor and is involved in B cell activation and differentiation. Activation of CD19 is accompanied by phosphorylation of the cytoplasmic domain, which promotes binding to kinases and the induction of intracellular signaling cascades. Mutations in CD19 can result in severe immunodeficiency syndromes. Clone 1D3 recognizes the same epitope as clone 6D5 in cross-competition assays.

Target Antigen Name: CD19
Alternative Names: B4
Gene ID: 12478
Species Reactivity: Mouse
Host Species: Rat (LEW)
Clonality: Monoclonal

Clone: 1D3

Isotype: IgG2a, kappa

Immunogen: Recombinant mouse CD19-transfected cell line

Conjugate: FITC

Applications

Verified: FC Reported: FC

Special Applications: This antibody clone has been verified for purity assessments of cells isolated with EasySep™ kits, including

EasySep™ Mouse B Cell Enrichment Kit (Catalog #19754).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.1% gelatin

Purification: The antibody was purified by affinity chromatography and conjugated with FITC under optimal conditions. The

solution is free of unconjugated FITC.

Stability and Storage: Product stable at 2 - 8°C when stored undiluted. Do not freeze, Protect product from prolonged exposure to

light. For product expiry date, please contact techsupport@stemcell.com.

Directions for Use: For flow cytometry the suggested use of this antibody is $\leq 0.25 \,\mu g$ per 1 x 10⁶ cells in 100 μL volume. It is


recommended that the antibody be titrated for optimal performance for each application.

Anti-Mouse CD19 Antibody, Clone 1D3, FITC

Antibodies



Data



Flow cytometry analysis of C57BL/6 mouse splenocytes labeled with Anti-Mouse CD19 Antibody, Clone 1D3, FITC (filled histogram) or a rat IgG2a, kappa isotype control antibody, FITC (solid line histogram).

Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

References

- 1. Von Muenchow L et al. (2014) The selection of mature B cells is critically dependent on the expression level of the co-receptor CD19. Immunol Lett 160(2): 113-19. (FC)
- 2. Chen Q et al. (2013) Retinoic acid and α-galactosylceramide regulate the expression of costimulatory receptors and transcription factors responsible for B cell activation and differentiation. Immunobiology 218(12): 1477-87. (FC, ICC, IF)
- 3. Yanaba K et al. (2013) CD19 expression in B cells regulates atopic dermatitis in a mouse model. Am J Pathol 182(6): 2214-22. (FC)
- 4. Ziegler AI et al. (2013) The CD19 signalling molecule is elevated in NOD mice and controls type 1 diabetes development. Diabetologia 56(12): 2659-68. (FC)
- 5. Keren Z et al. (2011) B-cell depletion reactivates B lymphopoiesis in the BM and rejuvenates the B lineage in aging. Blood 117(11): 3104-12. (FA, FC)
- 6. Depoil D et al. (2008) CD19 is essential for B cell activation by promoting B cell receptor-antigen microcluster formation in response to membrane-bound ligand. Nat Immunol 9(1): 63-72. (ICC, IF)
- 7. Shoham T et al. (2003) The tetraspanin CD81 regulates the expression of CD19 during B cell development in a postendoplasmic reticulum compartment. J Immunol 171(8): 4062-72. (FACS, FC, IP)
- 8. Carter RH et al. (2002) Role of CD19 signal transduction in B cell biology. Immunol Res 26(1-3): 45-54.
- 9. Gommerman JL et al. (2000) A role for CD21/CD35 and CD19 in responses to acute septic peritonitis: a potential mechanism for mast cell activation. J Immunol 165(12): 6915-21. (FC, ICC, IF)
- 10. Krop I et al. (1996) Self-renewal of B-1 lymphocytes is dependent on CD19. Eur J Immunol 26(1): 238-42. (FA, IP)
- 11. Krop I et al. (1996) The signaling activity of murine CD19 is regulated during B cell development. J Immunol 157(1): 48-56. (FC, IP)
- 12. Rickert RC et al. (1995) Impairment of T-cell-dependent B-cell responses and B-1 cell development in CD19-deficient mice. Nature 376(6538): 352-55.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485 MEDICAL DEVICE STANDARDS.

Copyright © 2015 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and EasySep are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. Alexa Fluor® is a registered trademark of Life Technologies Corporation. This product is licensed for internal research use only and its sale is expressly conditioned on the buyer not using it for manufacturing, performing a service, or medical test, or otherwise generating revenue. For use other than research, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.