

Cytokines

Human Recombinant BMP-2



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Bone morphogenetic protein 2

Catalog #	78004	50 µg
	78004.1	100 µg
	78004.2	1000 µg

Product Description

Bone morphogenetic protein 2 (BMP-2) is a member of the transforming growth factor beta (TGF- β) superfamily. BMP-2 is a disulfide-linked homodimer, acts as a ligand for complexes of type I and II BMP receptors, and primarily activates SMAD1/5/8 signaling (Nohe et al.). BMP-2 is a potent differentiation factor and directs human pluripotent stem cells (hPSCs) towards various cell types including extra-embryonic endoderm, mesenchymal cells, and chondrocytes (Pera et al.). Although BMP-2 expression is low in healthy cartilage, its expression is upregulated at the site of cartilage damage (Blaney Davidson et al.), and in vitro BMP-2 induces bone and cartilage formation and is able to induce chondrogenesis in human mesenchymal stem cells (Schmitt et al.).

Product Information

Alternative Names:	BDA2, BMP-2A, BMP2, Bone morphogenetic protein 2A
Accession Number:	P12643
Amino Acid Sequence:	MQAKHKQRKR LKSSCKRHPL YVDFSDVGWN DWIVAPPGYH AFYCHGECPF PLADHLNSTN HAIVQTLVNS VNSKIPKACC VPTLSAISM LYLDENEKVV LKNYQDMVVE GCGCR
Predicted Molecular Mass:	26 kDa
Species:	Human
Cross Reactivity:	Mouse, Rat, Monkey
Formulation:	Lyophilized after dialysis against acetic acid.
Source:	E. coli


Specifications

Activity:	The specific activity is $\geq 5 \times 10^3$ units/mg ($EC_{50} \leq 0.2$ µg/mL) as determined by the alkaline phosphatase activity induced in ATDC-5 cells.
Purity:	$\geq 95\%$
Endotoxin Level:	Measured by kinetic limulus amebocyte lysate (LAL) analysis and is ≤ 0.2 EU/µg protein.

Preparation and Storage

Storage:	Store at -80°C.
Stability:	Stable as supplied for 12 months from date of receipt.
Reconstitution:	Centrifuge vial before opening. Resuspend the product in sterile 20 mM acetic acid or 5 mM hydrochloric acid to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. Store at 2 - 8°C for up to 2 weeks or at -20°C to -80°C for up to 3 months. Avoid repeated freeze-thaw cycles.

Data



(A) The biological activity of Human Recombinant BMP-2 was tested by its ability to promote alkaline phosphatase production of ATDC-5 cells. Alkaline phosphatase production was measured using a fluorometric assay method. The EC₅₀ is defined as the effective concentration of the growth factor at which alkaline phosphatase production is at 50% of maximum. The EC₅₀ in the above example is 0.09 ng/mL.

(B) 2 µg of Human Recombinant BMP-2 was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human Recombinant BMP-2 has a predicted molecular mass of 26 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

- Blaney Davidson EN et al. (2007) Elevated extracellular matrix production and degradation upon bone morphogenetic protein-2 (BMP-2) stimulation point toward a role for BMP-2 in cartilage repair and remodeling. *Arthritis Res Ther* 9(5): R102.
- Nohe A et al. (2001) The Mode of Bone Morphogenetic Protein (BMP) Receptor Oligomerization Determines Different BMP-2 Signaling Pathways. *J Bio Chem* 277: 5330–8.
- Pera MF et al. (2004) Regulation of human embryonic stem cell differentiation by BMP-2 and its antagonist noggin. *J Cell Sci* 117(Pt 7): 1269–80.
- Schmitt B et al. (2003) BMP2 initiates chondrogenic lineage development of adult human mesenchymal stem cells in high-density culture. *Differentiation* 71(9-10): 567–77.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.