

**Anti-Mouse CD54 (ICAM-1)
Antibody, Clone YN1/1.7.4**



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Antibodies

Rat monoclonal IgG2b antibody
against mouse CD54, unconjugated

Catalog #60151
#60151.1

500 µg 0.5 mg/mL
50 µg 0.5 mg/mL

Product Description

The YN1/1.7.4 antibody reacts with CD54, a 90 - 110 kDa type I transmembrane glycoprotein that is a member of the Ig superfamily. CD54 consists of five Ig-like C2 type domains and has two isoforms generated from alternative splicing. As an important intercellular adhesion molecule, CD54 is involved in inflammatory reactions, antigen-specific immune responses, lymphocyte trafficking and T cell signal transduction events. CD54 mediates important cell-cell interactions and adhesion reactions by binding to its major ligands, LFA-1 (CD11a/CD18) and Mac-1 (CD11b/CD18). CD54 may also act as a receptor for rhinoviruses and some coxsackieviruses. CD54 is expressed on lymphocytes, monocytes, macrophages, dendritic cells, endothelial cells, and high endothelial venules. Expression of CD54 is upregulated on activated lymphocytes and also, following stimulation with inflammatory mediators such as cytokines, LPS and IFN-γ. The YN1/1.7.4 antibody has been employed to inhibit CD54-mediated functions (ie, cell-cell adhesion, antigen presentation to T cells and leukocyte migration) and to block the binding of CD54 to LFA-1 and Mac-1. The YN1/1.7.4 antibody recognizes a distinct epitope from that of KAT-1.

Target Antigen Name:	CD54
Alternative Names:	BB2, Cell surface glycoprotein P3.58, Human rhinovirus receptor, Intercellular adhesion molecule 1, Ly-47, MALA2, MyD10
Gene ID:	15894
Species Reactivity:	Mouse
Host Species:	Rat (Fisher 344)
Clonality:	Monoclonal
Clone:	YN1/1.7.4
Isotype:	IgG2b, kappa
Immunogen:	Mouse NS-1 myeloma cell line
Conjugate:	Unconjugated

Applications


Verified:	FC
Reported:	Block, ELISA, FC, IHC, IF, Inhibition, IP, Purification, RIA, WB

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Purification:	The antibody was purified by affinity chromatography.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. For product expiry date, please contact techsupport@stemcell.com .
Directions for Use:	The suggested use of this antibody is: FC, ≤ 0.25 µg per 1 × 10 ⁶ cells in 100 µL volume; WB, ≤ 4 µg/mL. It is recommended that the antibody be titrated for optimal performance for each application.

Data



Flow cytometry analysis of C57BL/6 mouse splenocytes labeled with Anti-Mouse CD54 (ICAM-1) Antibody, Clone YN1/1.7.4, followed by a mouse anti-rat IgG2b antibody, FITC (filled histogram) or Rat IgG2b, kappa Isotype Control Antibody, Clone RTK4530 (Catalog #60077), followed by a mouse anti-rat IgG2b antibody, FITC (solid line histogram).

Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

References

1. Zhang M et al. (2015) Real-time in vivo imaging reveals the ability of neutrophils to remove *Cryptococcus neoformans* directly from the brain vasculature. *J Leukoc Biol*. [Epub ahead of print] (IF, IHC)
2. Greineder CF et al. (2013) Vascular immunotargeting to endothelial determinant ICAM-1 enables optimal partnering of recombinant scFv-thrombomodulin fusion with endogenous cofactor. *PLoS One* 8(11): e80110. (IF, Live Cell ELISA, RIA)
3. Seidler DG et al. (2011) The role for decorin in delayed-type hypersensitivity. *J Immunol* 187(11): 6108–19. (WB)
4. Bankoti J et al. (2010) Effects of TCDD on the fate of naive dendritic cells. *Toxicol Sci* 115(2): 422–34. (FC)
5. Sumagin R & Sarelius IH. (2010) Intercellular adhesion molecule-1 enrichment near tricellular endothelial junctions is preferentially associated with leukocyte transmigration and signals for reorganization of these junctions to accommodate leukocyte passage. *J Immunol* 184(9): 5242–52. (Block, IF/Intravital Microscopy)
6. Sumagin R & Sarelius IH. (2007) A role for ICAM-1 in maintenance of leukocyte-endothelial cell rolling interactions in inflamed arterioles. *Am J Physiol Heart Circ Physiol* 293(5): H2786–98. (Block, IF/Intravital Microscopy)
7. Burns AR et al. (1994) Quantitation of ICAM-1 expression in mouse lung during pneumonia. *J Immunol* 153(7): 3189–98. (IHC)
8. Horley KJ et al. (1989) Molecular cloning of murine intercellular adhesion molecule (ICAM-1). *EMBO J* 8(10): 2889–96. (Inhibition, Purification)
9. Takei F. (1985) Inhibition of mixed lymphocyte response by a rat monoclonal antibody to a novel murine lymphocyte activation antigen (MALA-2). *J Immunol* 134(3): 1403–7. (Inhibition, IP)

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.