

Small Molecules

STAT5 Inhibitor

JAK/STAT pathway inhibitor; Inhibits STAT5

Catalog # 73852

5 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

STAT5 Inhibitor is a cell-permeable, nonpeptidic nicotinoyl hydrazone that specifically suppresses STAT5 ($IC_{50} = 47 \mu M$) via binding to its SH2 domain. It shows reduced potency towards the SH2 domains of STAT1, STAT3, or LCK (Müller et al.).

Molecular Name: STAT5 Inhibitor

Alternative Names: Not applicable

CAS Number: 285986-31-4


Chemical Formula: $C_{16}H_{11}N_3O_3$

Molecular Weight: 293.3 g/mol

Purity: $\geq 95\%$

Chemical Name: 2-[(4-oxo-4H-1-benzopyran-3-yl)methylene]hydrazide 3-pyridinecarboxylic acid

Structure:



Properties

Physical Appearance: A crystalline solid

Storage: Product stable at $-20^{\circ}C$ as supplied. Protect product from prolonged exposure to light. For long-term storage store with a desiccant. For product expiry date, please contact techsupport@stemcell.com.

Solubility: · DMSO ≤ 17 mM

For example, to prepare a 10 mM stock solution in DMSO, resuspend 1 mg in 341 μL of DMSO.

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at $-20^{\circ}C$. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

Published Applications

IMMUNOLOGY

- Induces partial apoptosis of mature eosinophils at high concentrations (Schwartz et al.).
- Rescues IL-17 production in a proportion of restimulated TH17 clones (Zielinski et al.).
- Reduces growth of induced regulatory T cells (iTregs) in vitro (Betts et al.).

References

- Betts BC et al. (2014) STAT5 polarization promotes iTregs and suppresses human T-cell alloresponses while preserving CTL capacity. *J Leukoc Biol* 95(2): 205–13.
- Müller J et al. (2008) Discovery of chromone-based inhibitors of the transcription factor STAT5. *Chembiochem* 9(5): 723–7.
- Schwartz C et al. (2015) Eosinophil-specific deletion of $\text{I}\kappa\text{B}\alpha$ in mice reveals a critical role of NF- κB -induced Bcl-xL for inhibition of apoptosis. *Blood* 125(25): 3896–904.
- Zielinski CE et al. (2012) Pathogen-induced human TH17 cells produce IFN- γ or IL-10 and are regulated by IL-1 β . *Nature* 484(7395): 514–8.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, please visit our website at www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.